

The inclusive $B \rightarrow \eta' X_s$ decay and $b \rightarrow sg^*$ form factors

Guey-Lin Lin[†]

Institute of Physics, National Chiao-Tung University, Hsin-chu 300, Taiwan

E-mail: glin@cc.nctu.edu.tw

Abstract

We compute the branching ratio of inclusive $B \rightarrow \eta' X_s$ decay based upon the QCD anomaly mechanism: $b \rightarrow s + g^* \rightarrow s + g + \eta'$. To obtain a reliable $B \rightarrow \eta' X_s$ branching ratio, we calculate the $b \rightarrow s + g^*$ form factors up to the next-to-leading-logarithmic(NLL) approximation. We point out that the Standard Model prediction for $B \rightarrow \eta' X_s$ is consistent with the CLEO data, in contrast to the claims of some previous works.

1. Introduction

The observations of exclusive $B \rightarrow \eta' K$ [1] and inclusive $B \rightarrow \eta' X_s$ [2] decays with high momentum η' mesons have stimulated many theoretical activities [3, 4, 5, 6, 7, 8, 9, 10]. The experimental fitting[2] shows that the dominant mechanism for the inclusive mode should be $b \rightarrow sg^* \rightarrow sg\eta'$ [3, 4] where the η' meson is produced via the anomalous $\eta' - g - g$ coupling. According to a previous analysis[4], this mechanism within the Standard Model(SM) can only account for 1/3 of the measured branching ratio: $\mathcal{B}(B \rightarrow \eta' X_s) = [6.2 \pm 1.6(\text{stat}) \pm 1.3(\text{syst})_{-1.5}^{+0.0}(\text{bkg})] \times 10^{-4}$ [2] with $2.0 < p_{\eta'} < 2.7$ GeV. Furthermore, the subleading mechanism for $B \rightarrow \eta' X_s$, based upon four-quark operators of the effective weak Hamiltonian[5, 6], is not sufficient to make up the above deficiency. Due to results of these analyses, the possibility of an enhanced $b \rightarrow sg$ or other mechanisms arising from physics beyond the Standard Model are raised to account for the above discrepancy in branching ratios[4, 6, 7]. In order to see if new physics should play any role in $B \rightarrow \eta' X_s$, one has to have a better understanding on the SM prediction. In this talk, we present a careful analysis on $B \rightarrow \eta' X_s$ [11] using the next-to-leading order effective Hamiltonian. In section 2, we illustrate how to compute the off-shell $b \rightarrow sg^*$ form factors in such a framework. In particular, the QCD equation of motion is applied to transform the charge-radius form factor of $b \rightarrow sg^*$ into the structures of certain four-quark operators. Therefore the effective weak Hamiltonian is shown applicable for computing such

a form factor. In section 3, we calculate the branching ratio and the recoil spectrum of $B \rightarrow \eta' X_s$ decay. The results are found to be consistent with the CLEO measurement[2]. Section 4 is the conclusion.

2. QCD equation of motion and $b \rightarrow s + g^*$ form factors

The effective Hamiltonian[12] relevant to the $B \rightarrow \eta' X_s$ decay is given by:

$$H_{eff} = -\frac{G_F}{\sqrt{2}} V_{ts}^* V_{tb} \left(\sum_{i=1}^6 C_i(\mu) O_i(\mu) + C_8(\mu) O_8(\mu) \right), \quad (1)$$

with

$$\begin{aligned} O_1 &= (\bar{s}_i c_j)_{V-A} (\bar{c}_j b_i)_{V-A}, \\ O_2 &= (\bar{s}_i c_i)_{V-A} (\bar{c}_j b_j)_{V-A}, \\ O_{3,5} &= (\bar{s}_i b_i)_{V-A} \sum_q (\bar{q}_j q_j)_{V \mp A}, \\ O_{4,6} &= (\bar{s}_i b_j)_{V-A} \sum_q (\bar{q}_j q_i)_{V \mp A}, \\ O_8 &= -\frac{g_s m_b}{4\pi^2} \bar{s}_i \sigma^{\mu\nu} P_R T_{ij}^a b_j G_{\mu\nu}^a, \end{aligned} \quad (2)$$

where $V \pm A \equiv 1 \pm \gamma_5$. Precisely speaking, this effective Hamiltonian can be used to calculate the off-shell $b \rightarrow sg^*$ form factors which are expressed as

$$\begin{aligned} \Gamma_\mu^{bsg} = & -\frac{G_F}{\sqrt{2}} V_{ts}^* V_{tb} \frac{g_s}{4\pi^2} (F_1 \bar{s}(q^2 \gamma_\mu - \not{q} q_\mu) L T^a b \\ & - i F_2 m_b \bar{s} \sigma_{\mu\nu} q^\nu R T^a b). \end{aligned} \quad (3)$$

[†] based upon a work done with X.-G. He

It is easily seen that $F_2 = -2C_8(\mu)$. However, the connection between F_1 and the effective Hamiltonian H_{eff} is less obvious. One may acquire some hints by rearranging the QCD penguin operators:

$$\begin{aligned} \sum_{i=3}^6 C_i O_i &= 2(C_4 + C_6)O_V - 2(C_4 - C_6)O_A \\ &+ (C_3 + \frac{C_4}{N_c})O_3 + (C_5 + \frac{C_6}{N_c})O_5, \end{aligned} \quad (4)$$

where

$$\begin{aligned} O_A &= \bar{s}\gamma_\mu(1 - \gamma_5)T^a b \sum_q \bar{q}\gamma^\mu\gamma_5 T^a q, \\ O_V &= \bar{s}\gamma_\mu(1 - \gamma_5)T^a b \sum_q \bar{q}\gamma^\mu T^a q. \end{aligned} \quad (5)$$

Since the light-quark bilinear in O_V carries the quantum number of a gluon, one expects[3] O_V give contributions to the $b \rightarrow sg^*$ form factors. In fact, by applying the QCD equation of motion, $D_\nu G_a^{\mu\nu} = g_s \sum_q \bar{q}\gamma^\mu T^a q$, we have $O_V = (1/g_s)\bar{s}\gamma_\mu(1 - \gamma_5)T^a b D_\nu G_a^{\mu\nu}$. In this form, O_V clearly contributes to the charge-radius form factor F_1 . Let us denote this part of contribution as F_1^a . We have $F_1^a = 4\pi/\alpha_s \cdot (C_4(\mu) + C_6(\mu))$. We remark that, at the NLL level, F_1 should also receive contributions from one-loop matrix elements. The dominant contribution, denoted as F_1^b , arises from the operator O_2 where its charm-quark-pair meets to form a gluon. In the NDR scheme, we find $F_1^b = 4\pi/\alpha_s \cdot (\bar{C}_4(q^2, \mu) + \bar{C}_6(q^2, \mu))$ where q^2 is the invariant mass of the gluon and

$$\begin{aligned} \bar{C}_4(q^2, \mu) &= \bar{C}_6(q^2, \mu) \\ &= \frac{\alpha_s(\mu)}{8\pi} C_2(\mu) \left(\frac{2}{3} + G(m_c^2, q^2, \mu) \right), \end{aligned} \quad (6)$$

with[13]

$$\begin{aligned} G(m_c^2, q^2, \mu) &= 4 \int_0^1 x(1-x)dx \\ &\times \log \left(\frac{m_c^2 - x(1-x)q^2}{\mu^2} \right). \end{aligned} \quad (7)$$

We point out that F_1^b , the O_2 contribution, is not negligible. For $\mu = 5$ GeV, one has $F_1^a = -4.03$ and $\text{Re}(F_1^b)$ (the real part of F_1^b) ranging between -1.5 and -3 for $0.2 < q^2/m_b^2 < 0.7$. The peak value $\text{Re}(F_1^b) \equiv -3$ occurs at the charm-pair threshold $q^2 = 4m_c^2$. For $q^2 > 4m_c^2$, F_1^b develops an imaginary part whose value is roughly $2i$. Concerning previous results on F_1 , we note that Ref. [3] intends to compute F_1 with effective weak

Hamiltonian. However, only the contribution by F_1^a is considered. Ref. [4] took $F_1 = -5$ which is a result of an one-loop calculation[15]. Clearly the q^2 dependencies of F_1 are also absent. As we shall see in the next section, the contribution by F_1^b , which is not included in previous works, can significantly enhance the $B \rightarrow \eta' X_s$ branching ratio such that SM prediction is consistent with the CLEO measurement.

3. The inclusive $B \rightarrow \eta' X_s$ decay

In this decay, the η' final state arises from the off-shell gluon splitting, $g^* \rightarrow g\eta'$, where g^* is produced through $b \rightarrow sg^*$. The branching-ratio distribution of $b(p) \rightarrow s(p') + g(k) + \eta'(k')$ is[4]:

$$\begin{aligned} \frac{d^2 \mathcal{B}(b \rightarrow sg\eta')}{dx dy} &\cong 0.2 \cos^2 \theta \left(\frac{g_s(\mu)}{4\pi^2} \right)^2 \frac{a_g^2(\mu) m_b^2}{4} \\ &\times \left(|\Delta F_1|^2 c_0 + \text{Re}(\Delta F_1 F_2^*) \frac{c_1}{y} \right. \\ &\left. + |\Delta F_2|^2 \frac{c_2}{y^2} \right), \end{aligned} \quad (8)$$

where $a_g(\mu) \equiv \sqrt{N_F} \alpha_s(\mu) / \pi f_{\eta'}$ is the strength of $\eta' - g - g$ vertex: $a_g \cos \theta \epsilon_{\mu\nu\alpha\beta} q^\alpha k^\beta$ with q and k the momenta of two gluons; $x \equiv (p' + k)^2/m_b^2$ and $y \equiv (k + k')^2/m_b^2$; c_0 , c_1 and c_2 are functions of x and y given by:

$$\begin{aligned} c_0 &= \left[-x^2 y + (1-y)(y-x')(x + \frac{y}{2} - \frac{x'}{2}) \right], \\ c_2 &= \left[x^2 y^2 - (1-y)(y-x')(xy - \frac{y}{2} + \frac{x'}{2}) \right], \\ c_1 &= (1-y)(y-x')^2, \end{aligned} \quad (9)$$

with $x' \equiv m_{\eta'}^2/m_b^2$; and the $\eta' - \eta$ mixing angle θ is taken to be -15.4° [16]. Following the approach in Ref. [4], we evaluate the $\alpha_s(\mu)$ in a_g at the running scale $\mu^2 = q^2$. Taking $\mu = 5$ GeV for evaluating other scale-dependent quantities, we find $\mathcal{B}(b \rightarrow sg\eta') = 6.4 \times 10^{-4}$ with the cut $m_X \equiv \sqrt{(k+p')^2} \leq 2.35$ GeV imposed in the CLEO measurement[2]. This branching ratio is consistent with CLEO's measurement on $\mathcal{B}(B \rightarrow \eta' X_s)$ branching ratio[2]. Without the kinematic cut, we obtain $\mathcal{B}(b \rightarrow sg\eta') = 1.2 \times 10^{-3}$, which is much larger than 4.3×10^{-4} calculated previously[4]. Clearly this enhancement is due to the contribution of F_1^b , which increases the magnitude of F_1 and thus enhances the the branching ratio of $b \rightarrow sg\eta'$ according to Eq. (8). Since Ref.[3] also neglects the contribution by F_1^b , its prediction on $\mathcal{B}(b \rightarrow sg\eta')$

would be much smaller than ours if the $\eta' - g - g$ coupling there is also evaluated at the running scale $\mu^2 = q^2$. However, the prediction by Ref. [3] is comparable to ours, since, in that work, the $\alpha_s(\mu)$ in a_g is evaluated at the lowest possible scale $\mu^2 = m_{\eta'}^2$, and the interference between the contributions by F_1 and F_2 is constructive rather than destructive reported here and in Ref.[4].

To ascertain our calculation, we also check the μ dependence of the $b \rightarrow sg\eta'$ branching ratio. Using NDR scheme with $\mu = 2.5$ GeV and imposing the kinematic cut $m_X \leq 2.35$ GeV, we find $\mathcal{B}(b \rightarrow sg\eta') = 7.1 \times 10^{-4}$, which is only 10% larger than the branching ratio obtained at $\mu = 5$ GeV. This insensitivity to the scale-choice is reassuring. We also obtain the spectrum $d\mathcal{B}(b \rightarrow sg\eta')/dm_X$ which has been shown in Ref.[11] and will not be displayed here. The peak of the spectrum corresponds to $m_X \approx 2.4$ GeV. As pointed out in Ref. [2], this type of spectrum gives the best fit to the CLEO data.

4. Concluding remarks

We have calculated the branching ratio of $b \rightarrow sg\eta'$ by including the NLL correction to the $b \rightarrow sg^*$ form factors. By evaluating the $\eta' - g - g$ coupling at the running scale $\mu = q^2$ and cutting the recoil-mass m_X at 2.35 GeV, we obtained $\mathcal{B}(b \rightarrow sg\eta') = (6.4 - 7.1) \times 10^{-4}$ depending on the choice of the renormalization scale for evaluating the $b \rightarrow sg^*$ form factors. We have not addressed the possible form-factor suppression of the $\eta' - g - g$ vertex, which occurs as the gluons attached to the vertex go farther off-shell[3, 4, 6]. So far it remains unclear how much the form-factor suppression might be. However, comparing our prediction with the CLEO measurement on $\mathcal{B}(B \rightarrow \eta' X_s)$, which still has a large error bar, it remains possible that the anomaly-induced process $b \rightarrow sg\eta'$ could account for the CLEO data within the framework of the Standard Model.

5. Acknowledgments

This work is supported by National Science Council of R.O.C. under the grant number NSC 87-2112-M-009-038, and National Center for Theoretical Sciences of R.O.C. under the topical program: PQCD, B and CP.

References

- [1] CLEO collaboration, B.H. Behrens *et al.* 1998 *Phys. Rev. Lett.* **80** 3710.
- [2] CLEO Collaboration T. E. Browder *et al.* 1998 *Phys. Rev. Lett.* **81** 1786.
- [3] D. Atwood and A. Soni 1997 *Phys. Lett. B* **405** 150.
- [4] W.S. Hou and B. Tseng 1998 *Phys. Rev. Lett.* **80** 434.
- [5] A. Datta, X.-G. He, and S. Pakvasa 1998 *Phys. Lett. B* **419** 369.
- [6] A. L. Kagan and A. Petrov, Preprint hep-ph/9707354; Preprint hep-ph/9806266.
- [7] H. Fritzsch 1997 *Phys. Lett. B* **415** 83; X.-G. He, W.-S. Hou and C.S. Huang 1998 *Phys. Lett. B* **429** 99.
- [8] H.-Y. Cheng, and B. Tseng 1998 *Phys. Rev. D* **58** 094005; A. Ali, J. Chay, C. Greub and P. Ko 1998 *Phys. Lett. B* **424** 161; N. Deshpande, B. Dutta and S. Oh 1998 *Phys. Rev. D* **57** 5723.
- [9] A. Ali, G. Kramer and C.-D. Lu 1998 *Phys. Rev. D* **58** 094009.
- [10] I. Halperin and A. Zhitnitsky 1998 *Phys. Rev. Lett.* **80** 438; F. Araki, M. Musakonov and H. Toki 1999 *Phys. Rev. D* **59** 037501; D.S. Du, Y.-D. Yang and G.-H. Zhu 1999 *Phys. Rev. Lett.* **82** 014007; M. Ahmady, E. Kou and A. Sugamoto 1998 *Phys. Rev. D* **58** 014015; D.S. Du, C.S. Kim and Y.-D. Yang 1998 *Phys. Lett. B* **426** 133; F. Yuan and K.-T. Chao 1997 *Phys. Rev. D* **56** 2495; A. Dighe, M. Gronau and J. Rosner 1997 *Phys. Rev. Lett.* **79** 4333.
- [11] X.-G. He and G.-L. Lin 1999 *Phys. Lett. B* **454** 123.
- [12] For an extensive review on the subject of effective Hamiltonian, see G. Buchalla, A. J. Buras and M. E. Lautenbacher 1996 *Review of Modern Physics* **68** 1125, which contains a detailed list of original literatures.
- [13] In the original article[11], we use the scheme-independent Wilson coefficient[14]. Hence the contribution of one-loop matrix element is also distinct. In particular, the constant added to $G(m_c^2, q^2, \mu)$ in Eq. (6) is $\frac{10}{9}$ rather than $\frac{2}{3}$. For detail, we refer the readers to Eqs. (4) and (5) of Ref.[11]. Note that the r.h.s. of Eq. (4) in Ref.[11] should be $P_s(q^2, \mu)$. The extra minus sign is a typing error.
- [14] A. Buras, M. Jamin, M. Lautenbacher and P. Weisz 1992 *Nucl. Phys. B* **370** 69; N. G. Deshpande and X.-G. He 1994 *Phys. Lett. B* **336** 471.
- [15] W. S. Hou, A. Soni and H. Steger 1987 *Phys. Rev. Lett.* **59** 1521.
- [16] T. Feldmann, P. Kroll and B. Stech 1998 *Phys. Rev. D* **58** 114006.